

**Table 29. Tuberculosis Cases by Initial Drug Regimen: 59 Reporting Areas, 1999**

Reporting Area	Total Cases	Cases Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percent of Cases with Initial Drug Regimen <sup>1</sup>		
			No.	%	IR	IRZ	IRZ,E/S <sup>2</sup>
<b>United States</b>	<b>17,531</b>	<b>16,998</b>	<b>16,919</b>	<b>99.5</b>	<b>2.2</b>	<b>11.2</b>	<b>76.8</b>
Alabama	314	301	301	100.0	2.7	70.1	23.3
Alaska	61	60	60	100.0	0.0	5.0	90.0
Arizona	262	250	250	100.0	0.0	9.6	79.2
Arkansas	181	175	171	97.7	43.9	37.4	14.6
California	3,606	3,513	3,506	99.8	1.3	6.2	85.1
Colorado	88	86	85	98.8	0.0	1.2	92.9
Connecticut	121	120	120	100.0	0.8	14.2	80.0
Delaware	34	34	34	100.0	0.0	8.8	88.2
District of Columbia	70	68	68	100.0	0.0	4.4	85.3
Florida	1,277	1,232	1,232	100.0	1.4	8.5	79.6
Georgia	665	646	642	99.4	0.9	13.6	77.1
Hawaii	184	178	176	98.9	7.4	16.5	71.0
Idaho	16	16	16	100.0	0.0	25.0	75.0
Illinois	825	808	797	98.6	1.5	7.7	73.4
Indiana	150	146	146	100.0	2.7	19.9	63.7
Iowa	58	57	57	100.0	1.8	21.1	66.7
Kansas	69	65	63	96.9	4.8	22.2	61.9
Kentucky	209	200	200	100.0	1.0	11.0	77.0
Louisiana	357	343	332	96.8	2.4	14.5	77.4
Maine	23	23	23	100.0	4.3	17.4	73.9
Maryland	294	286	286	100.0	1.0	6.6	87.8
Massachusetts	270	264	264	100.0	0.8	6.1	84.1
Michigan	351	337	337	100.0	5.0	32.9	51.6
Minnesota	201	198	198	100.0	0.5	16.7	71.7
Mississippi	215	208	207	99.5	1.4	16.4	79.7
Missouri	208	202	201	99.5	2.0	8.5	72.6
Montana	14	13	13	100.0	0.0	15.4	84.6
Nebraska	18	18	18	100.0	11.1	5.6	61.1
Nevada	93	92	92	100.0	1.1	5.4	87.0
New Hampshire	19	18	18	100.0	0.0	22.2	77.8
New Jersey	571	548	548	100.0	2.0	9.3	73.9
New Mexico	64	57	57	100.0	1.8	33.3	61.4
New York State <sup>3</sup>	377	364	364	100.0	1.6	8.0	81.3
New York City	1,460	1,426	1,426	100.0	1.1	3.6	81.6
North Carolina	488	477	476	99.8	0.4	3.8	89.3
North Dakota	7	7	7	100.0	14.3	14.3	71.4
Ohio	317	305	304	99.7	5.9	22.7	56.9
Oklahoma	208	199	199	100.0	8.5	17.1	58.8
Oregon	123	122	118	96.7	0.8	4.2	90.7
Pennsylvania	454	432	417	96.5	2.6	9.4	78.4
Rhode Island	53	53	52	98.1	3.8	1.9	84.6
South Carolina	315	302	302	100.0	1.0	16.2	64.2
South Dakota	21	19	19	100.0	0.0	10.5	78.9
Tennessee	382	371	371	100.0	2.4	19.4	68.5
Texas	1,649	1,598	1,586	99.2	1.8	10.2	77.7
Utah	40	36	36	100.0	5.6	16.7	69.4
Vermont	3	3	3	100.0	0.0	0.0	66.7
Virginia	334	324	324	100.0	1.2	14.8	77.5
Washington	258	250	250	100.0	1.6	6.0	89.6
West Virginia	41	39	39	100.0	10.3	35.9	35.9
Wisconsin	110	106	105	99.1	3.8	9.5	75.2
Wyoming	3	3	3	100.0	0.0	0.0	66.7
American Samoa <sup>4</sup>	4	4	4	100.0	0.0	0.0	100.0
Fed. States of Micronesia <sup>4</sup>	...	...	...	...	...	...	...
Guam <sup>4</sup>	69	67	67	100.0	0.0	0.0	86.6
N. Mariana Islands <sup>4</sup>	66	64	64	100.0	0.0	4.7	90.6
Puerto Rico <sup>4</sup>	200	181	181	100.0	0.0	1.7	93.4
Republic of Palau <sup>4</sup>	11	11	11	100.0	9.1	9.1	63.6
U.S. Virgin Islands <sup>4</sup>	...	...	...	...	...	...	...

1. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for >75% of cases.

2. I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

3. Excludes New York City.

4. Not included in U.S. totals.

Ellipses indicate data not available.

**Note:** Excluding cases with no information on initial drug regimen, 159 (0.9%) were not started on any drugs, 27 (0.2%) were started on one drug, and 1,463 (8.6%) had an initial multidrug regimen other than IR, IRZ, and IRZ,E/S.