

Table 5. Tuberculosis Cases by Case Verification Criterion and by Site of Disease: United States, 1989-1999

Year	Total Cases	Verification Criterion ¹								Site of Disease			
		Positive Culture		Positive Smear		Clinical Case Definition		Provider Diagnosis		Pulmonary ²		Extra-pulmonary	
		No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1989	23,495	19,483	83	351	1	2,475	11	1,186	5	19,639	84	3,835	16
1990	25,701	20,897	81	556	2	2,958	12	1,290	5	21,576	84	4,091	16
1991	26,283	21,417	81	388	1	2,992	11	1,486	6	21,937	83	4,327	16
1992	26,673	21,398	80	407	2	3,141	12	1,727	6	22,371	84	4,288	16
1993	25,287	20,081	79	309	1	2,994	12	1,903	8	21,255	84	3,995	16
1994	24,361	19,537	80	236	1	2,794	11	1,794	7	20,385	84	3,964	16
1995	22,860	18,292	80	220	1	2,664	12	1,684	7	18,991	83	3,860	17
1996	21,337	17,234	81	150	1	2,556	12	1,397	7	17,445	82	3,870	18
1997	19,851	16,015	81	177	1	2,355	12	1,304	7	16,285	82	3,554	18
1998	18,361	14,830	81	166	1	2,207	12	1,158	6	14,813	81	3,541	19
1999	17,531	13,997	80	176	1	2,058	12	1,300	7	14,083	80	3,438	20

1. Based on the public health surveillance case definition for tuberculosis: CDC. Case definitions for infectious conditions under public health surveillance. MMWR 1997;46(No. RR-10):40-41. See Appendix B.

2. Includes cases of both pulmonary and extrapulmonary disease and cases of miliary TB.

Note: See Technical Notes (Appendix A) for a description of national TB surveillance.

Table 6. Pulmonary Tuberculosis Cases by Sputum Smear and Sputum Culture Results: United States, 1989-1999

Year	Total Pulmonary Cases ¹	Sputum Smear Results						Sputum Culture Results					
		Positive		Negative		Not Done or Unknown		Positive		Negative		Not Done or Unknown	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1989	19,639	8,759	(45)	5,924	(30)	4,956	(25)	13,712	(70)	1,908	(10)	4,019	(20)
1990	21,576	9,391	(44)	6,865	(32)	5,320	(25)	14,816	(69)	2,124	(10)	4,636	(21)
1991	21,937	9,095	(41)	7,281	(33)	5,561	(25)	15,022	(68)	2,232	(10)	4,683	(21)
1992	22,371	8,975	(40)	7,413	(33)	5,983	(27)	15,124	(68)	2,476	(11)	4,771	(21)
1993	21,255	9,324	(44)	7,747	(36)	4,184	(20)	14,708	(69)	2,675	(13)	3,872	(18)
1994	20,385	8,845	(43)	7,770	(38)	3,770	(18)	14,080	(69)	2,618	(13)	3,687	(18)
1995	18,991	8,068	(42)	7,717	(41)	3,206	(17)	13,236	(70)	2,597	(14)	3,158	(17)
1996	17,445	7,449	(43)	7,337	(42)	2,659	(15)	12,232	(70)	2,507	(14)	2,706	(16)
1997	16,285	6,882	(42)	6,878	(42)	2,525	(16)	11,481	(71)	2,226	(14)	2,578	(16)
1998	14,813	6,630	(45)	6,016	(41)	2,167	(15)	10,472	(71)	2,101	(14)	2,240	(15)
1999	14,083	6,252	(44)	5,626	(40)	2,205	(16)	9,777	(69)	2,049	(15)	2,257	(16)

1. Includes cases of both pulmonary and extrapulmonary disease and cases of miliary TB.