

# Introduction to Radiology For TB Nurses

Megan Devine, MD

Medical Consultant Heartland National TB Center


Associate Professor Internal Medicine/Pulmonary Medicine/Critical Care University of Texas Health Science Center at Tyler


# Chest Radiology in TB

- X-Rays penetrate air, absorbed by fluids/solids
- Chest radiographs show us shadows of organs and structures in the chest
- Interpretation of a chest radiograph is pattern recognition that requires clinical correlation for true diagnosis
- To know what "abnormal" looks like, you have to know what "normal" looks like

# Chest Radiology Basics

- Dark/black = Air
- Dense White = Calcium (Bone Density)
- White = Water Density (Everything else)
  - Water
  - Blood
  - Fat
  - Tissue
  - Pus





# Lymph nodes in the chest

1 Low cervical, supraclavicular and sternal notch nodes

2R Upper Paratracheal (right)

4R Lower Paratracheal (right)

4L Lower Paratracheal (left)

5 Subaortic

Subcarinal 7


Paraesophageal (below carina) 8

9

10 Hilar

11 Interlobar

12 Lobar




#### Normal CXR Child



#### Role of CT in the Diagnosis of TB

- CT is <u>not</u> the primary radiologic diagnostic test for TB (CT is overused in the US)
- Usually don't need CT for cavitary consolidation
- If TB is a possible diagnosis, sputum for AFB should be obtained prior to CT
- In most instances, CT should be reserved for patients in whom the diagnosis is unclear





**Onward Transmission** 


Cellestis


### Primary Tuberculosis

- Most commonly in children and immune compromised patients
- Opacities are seen the in middle and lower lungs; commonly unilateral, bilateral 15%
- Hilar or paratracheal lymphadenopathy with or without infiltrates is characteristic.
- Lymph node enlargement may cause bronchial compression
- Pleural effusion (25% can occur in primary disease)













#### Post Primary, Reactivation Tuberculosis

- Characterized by upper lobe predilection, cavitation and absence of lymphadenopathy.
- Cavitation is the hallmark; can also see parenchymal disease (consolidation), hematogenous dissemination (milliary), bronchogenic spread (tree-in-bud) and pleural disease.
- Fibrosis and calcification are seen after healing.





#### Millet Seeds

Slender plant, 1-15 feet

Seeds ~ 2-3 mm in diameter

Africa and India














### Tree in Bud.....









## Pleural Effusions


- Primary TB (25%)
- Hypersensitivity reaction to TB proteins
- Organisms uncommonly isolated from fluid
- May not be associated with obvious parenchymal disease on CXR








# Back to our Eritrean family...



#### May 2019

37 year old African man 4 months of cough, weight loss, and poor energy

### Chest xray is one piece of the TB puzzle





Sputum AFB smear and PCR +, culture + MTB

# Chest Radiology in TB

 Interpretation of a chest radiograph is pattern recognition that requires clinical correlation for true diagnosis

# Thank you

Heartland National TB Center

