



Developing a TB Patient Care Plan

Jacqueline I Maldonado, RN

July 07, 2022

Introduction to TB Nurse Case Management Online

July 2022

Jacqueline I Maldonado, RN has the following disclosures to make:

No conflict of interest

No relevant financial relationships with any commercial companies pertaining to this educational activity



What is a Care Plan?


1. A formal process that

- provides direction
- helps to identify existing needs
- recognize potential needs or risks.

2. Main focus

- facilitate standardized, evidence-based and holistic care.

3. Provide a means of communication to achieve health care outcomes.



Objective: Have TB Nurse Case Manager develop a TB Patient Care Plan

- By understanding the following:
 - ✓ Description of treatment regimen
 - ✓ Methods of monitoring for adverse reactions
 - ✓ Methods of assessing and ensuring adherence to treatment
 - ✓ Methods for evaluating treatment response



TB Treatment Goals:

1. Cure the patient
2. Prevent death, disability or drug resistance
3. Prevent further transmission



There is no “i” in team, but there is in responsible.



Patient-Centered Care

“Patient-centered care is providing care that is respectful of and responsive to individual patient preferences, needs and values, and ensuring that patient values guide all clinical decisions.”

- Institute of Medicine (IOM)



Patient Education Topics:

- TB Disease Process
- What medication should be taken, how much and how often
- Possible adverse reactions to the medications
- When to seek necessary medical attention
- Consequences of not taking their medicine correctly
- TB infection, restriction measures, and isolation precautions



Understand Your Patient

- Patient's perception & knowledge of TB
- Sociocultural influences
- Home and work habits
- Patient's support system



TREATMENT REGIMEN



Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis

Payam Nahid,¹ Susan E. Dorman,² Narges Alipanah,¹ Pennan M. Barry,³ Jan L. Brozek,⁴ Adithya Cattamanchi,¹ Lelia H. Chaisson,¹ Richard E. Chaisson,² Charles L. Daley,⁵ Malgosia Grzemska,⁶ Julie M. Higashi,⁷ Christine S. Ho,⁸ Philip C. Hopewell,¹ Salmaan A. Keshavjee,⁹ Christian Lienhardt,⁶ Richard Menzies,¹⁰ Cynthia Merrifield,¹ Masahiro Narita,¹² Rick O'Brien,¹³ Charles A. Peloquin,¹⁴ Ann Raftery,¹ Jussi Saukkonen,¹⁵ H. Simon Schaaf,¹⁶ Giovanni Sotgiu,¹⁷ Jeffrey R. Starke,¹⁸ Giovanni Battista Migliori,¹¹ and Andrew Vernon⁸

¹University of California, San Francisco; ²Johns Hopkins University, Baltimore, Maryland; ³California Department of Public Health, Richmond; ⁴McMaster University, Hamilton, Ontario, Canada; ⁵National Jewish Health, Denver, Colorado; ⁶World Health Organization, Geneva, Switzerland; ⁷Tuberculosis Control Section, San Francisco Department of Public Health, California; ⁸Division of Tuberculosis Elimination, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention, Atlanta, Georgia; ⁹Harvard Medical School, Boston, Massachusetts; ¹⁰McGill University, Montreal, Quebec, Canada; ¹¹WHO Collaborating Centre for TB and Lung Diseases, Fondazione S. Maugeri Care and Research Institute, Tradate, Italy; ¹²Tuberculosis Control Program, Seattle and King County Public Health, and University of Washington, Seattle; ¹³Ethics Advisory Group, International Union Against TB and Lung Disease, Paris, France; ¹⁴University of Florida, Gainesville; ¹⁵Durban University, Durban, South Africa; ¹⁶Department of Biomedicine and Child Health, Stellenbosch University, Cape Town, South Africa; ¹⁷University of Perugia, Italy

CDC Recommendation for Treatment of Drug-Susceptible TB is a 4 drug regimen, aka **RIPE:**

Rifampin – RIF

Isoniazid – INH

Pyrazinamide – PZA

Ethambutol - EMB

Doses are based on weight and age of patient

Dosing Recommendations for Adult Patients with Drug-Susceptible Organisms

Drug	Normal Renal Function	Change in Frequency for Reduced Renal Function?	Creatinine Clearance <30 mL/min*
Ethambutol	<u>Standard dose^b:</u> 15-20 mg/kg once daily	Yes	20-25 mg/kg 3x/weekly (not daily)
Isoniazid	<u>Standard dose:</u> 5 mg/kg daily (max 300 mg) Vitamin B6 daily 25-50 mg <u>Intermittent dose:</u> 15 mg/kg (max 900 mg) <u>High dose therapy:</u> 13-18 mg/kg daily	No	No dose adjustment
Pyrazinamide	<u>Standard dose^b:</u> 25-35 mg/kg daily	Yes	25-35 mg/kg (maximum 3000 mg) 3x/weekly (not daily)
Rifabutin	<u>Standard dose:</u> 300 mg daily	No	Monitor drug concentrations to avoid toxicity
Rifampin	<u>Standard dose:</u> 10 mg/kg daily	No	No dose adjustment

*Including adult patients receiving hemodialysis

^bBased on estimated lean body weight. Optimal doses for obese patients are not established.

Please note: Standard doses are given unless there is intolerance; there should be careful monitoring of neurotoxicity; the medications should be given after hemodialysis on the day of hemodialysis; and monitoring of serum drug concentrations should be considered.



Tuberculosis Treatment Guidelines

Drug Regimens for Microbiologically Confirmed Pulmonary Tuberculosis Caused by Drug-Susceptible Organisms

Dosing Recommendations for Adult Patients with Drug-Susceptible Organisms

Adapted from the Official American Thoracic Society, Centers for Disease Control and Prevention, Infectious Disease Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis Clinical Infectious Diseases • 2016

This publication was supported by the Grant or Cooperative Agreement Number NU52PS910161 funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.

2303 Southeast Military Drive • San Antonio, Texas 78223

1-800-TEX-LUNG • www.heartlandnbtbc.org

2 Phases of Treatment

Initial Phase + Continuation Phase = Length of Treatment

8 weeks + 18 weeks = 6 months of treatment

8 weeks + 26 weeks = 9 months of treatment



Initial Phase of Treatment

RIPE + B6

24 doses if receiving 3 times/week

40 doses if receiving 5 days/week

56 doses if receiving 7 days/week

8 weeks

This phase must be completed before proceeding to the continuation phase



Continuation Phase of Treatment

RIF + INH + B6

54 doses if receiving 3 days/week

90 doses if receiving 5 days/week

126 doses if receiving 7 days/weeks

18 weeks versus 26 weeks



Why Directly Observed Therapy (DOT)?

1. Provides visual evaluation/observation of patient tolerance of medication
2. Provides visual evaluation/observation of patient response to treatment
3. Provides daily opportunity for patient education



Side Effect vs. Adverse Drug Reaction (ADR)

Side Effect:

- a less precise term, often refers to milder, predictable effects of taking a medication.
- Examples:
 - Discolored body fluids from Rifampin
 - Decrease effectiveness of birth control pills/implants from Rifampin

Adverse Drug Reaction (ADR):

- As defined by the World Health Organization (WHO), an Adverse Drug Reaction is a response to a drug that is noxious and unintended and occurs at doses normally used in man for the prophylaxis, diagnosis or therapy of a disease.
- Examples:
 - Hepatitis
 - Rash

MONITORING FOR ADVERSE REACTION

- Patient education
- Daily by DOT Provider
 - Observation
 - Toxicity screen
- Monthly or as ordered by licensed healthcare worker
 - Observation
 - Toxicity assessment
 - Labs



Blood Analysis:

- Baseline
- Monthly
- Complaint or adverse reaction
- Special situations



ADHERENCE TO TB TREATMENT

- Assessing Adherence > daily CM's duty
 - Designated place and time for DOT
 - Appointments met and rescheduled
- Ensuring Adherence > possible challenge
 - Incentives (rewards given to patients to encourage taking DOT or attend clinic appointments, such as food, clothing or personal products)
 - Enablers (ex. helps patient receive treatment, such as transportation vouchers to get to the clinic, appointment reminders and social service assistance)



EVALUATING TREATMENT RESPONSE

- Clinical
- Bacteriological
- Radiographic



Clinical

- Medication tolerance
- TB symptom improvement
- Appetite status
- Activity level
- Affect – mental status



- 

Genotype Number: _____

[illegible]

Radiographic Imaging

- After 2 months of TB medication
 - CXR or CT (depending on site of disease)
 - Improved from baseline?
- At end of TB treatment
 - CXR or CT (depending on site of disease)
 - Improved from previous image
- As needed based on patient findings



Indicators of poor response to Treatment

- Clinically - No improvement
- Bacteriologically – minimal to no improvement
- Radiologically – no improvement or worsening



Reasons for Poor Response

- Poor DOT adherence
- Patient vomiting after taking TB medication
- Poor absorption of medications
- Development of Drug Resistance
- Patient “Cheeking” Pills



Patient Centered Care

- Build a relationship with your patient
- Educate your patient and their family
- Provide the Right drugs, Right dosage, and Right number of doses
- Perform Baseline evaluations
- Use logs and graphs to monitor progress
- Document and communicate with the TB team
- Screen regularly for medication side effects and adverse reactions.
- Evaluate for improvement clinically and diagnostically



Questions?



References

- American Thoracic Society, Center for Disease Control and Prevention, Infectious Diseases Society of America Treatment of tuberculosis. MMWR 2003.
- Brenner, Ben. (2017). What is Patient Centered Care and How to Provide It. MedPro Disposal. <https://www.medprodisposal.com/practice-management/patient-centered-care-provide/>
- Center for Disease Control. (2014). Core Curriculum Chapter 6: Treatment of tuberculosis disease. Division of Tuberculosis Elimination. <https://www.cdc.gov/tb/education/corecurr/pdf/chapter6.pdf>
- Center for Disease Control. Controlling tuberculosis in the United States: Recommendations from the American Thoracic Society, CDC, and the Infectious Diseases Society of America. MMWR 2005; 54 (No. RR-12). www.cdc.gov/mmwr/preview/mmwrhtml/rr5412a1.htm
- Infectious Disease Society of America. (2016). Official American Thoracic Society/Infectious Diseases Society of America/Centers for Disease Control and Prevention Clinical Practice Guidelines: Diagnosis of Tuberculosis in Adults and Children. <https://www.thoracic.org/statements/resources/tb-opi/diagnosis-of-tuberculosis-in-adults-and-children.PDF>
- Treatment of Tuberculosis: Guidelines. 4th edition. Geneva: World Health Organization; 2010. 6, Supervision and patient support. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK138737/>
- Tuberculosis Nursing: A Comprehensive Guide to Patient Care, 2nd Edition, v. 06/13/11.

