

Importance of Weight in the Treatment Outcomes of a Patient with TB


Essentials of Nurse Case Management

Catalina B. Navarro BSN, RN
Nurse Consultant/Educator

1

Objectives

- Discuss the Importance of weight gain on TB treatment outcomes
- Demonstrate the use of the BMI chart with case studies



HEARTland
NATIONAL TB CENTER
Tuberculosis in Texas, Texas, and the World

Impact of Poor Nutrition on TB Relapse

Reference: Lack of Weight Gain and Relapse Risk in a Large Tuberculosis Treatment Trial: Awa Khan, Timothy R. Sterling, Randall Reeves, Andrew Vernon, C. Robert Horsburgh and the Tuberculosis Trials Consortium; American Journal of Respiratory and Critical Care Medicine Vol 174, pp. 344-348, (2006)

Weight gain of 5% or less during the first 2 months of therapy is associated with an increased risk of relapse, even after controlling for other factors.

Consultation to healthcare providers at 1-800-TEX-LUNG
2303 SE Military Drive, San Antonio, TX 78223
www.HeartlandNTBC.org

This product produced with funds awarded by the Centers for Disease Control & Prevention (CDC)
Created 3/08

CHART 1. BODY MASS INDEX (BMI)

Legend: Underweight (blue), Weight Appropriate (green), Overweight (yellow), Obese (red)

2

Malnutrition


Malnutrition refers to **deficiencies, excesses, or imbalances** in a person's intake of energy and/or nutrients.

- **Undernutrition**
- Micronutrient-related malnutrition
- Overweight and obesity




3

Geographic Overlap between TB and Undernutrition Worldwide 2018



4

Geographic Overlap between TB and Undernutrition Worldwide 2018



5

For a TB-free India, break the cycle of hunger and disease

Malnutrition and tuberculosis are India's major public health challenges. And the importance of nutritional intervention as a weapon against tuberculosis cannot be overemphasised

ANALYSIS Updated: Sep 08, 2017 17:48 IST

ht Rajan Sankar



People suffering from undernutrition are predisposed to contracting TB. In India, undernutrition contributes to a staggering 55% of the annual TB incidence.

6

Undernutrition and TB

"Rise in tuberculosis mortality was recorded in 1914-1916, and in those years the consumption of bread and flour rose, **whereas that of meat decreased.**"--

"High TB mortality in Europe during and since WWII, coincided with **great reduction of intake of protein food**, such as, meat, fish and eggs"

[Sandler MD \(Diet Prevents Polio\)](#)

7

TB Incidence Related to BMI 1971-1992




Figure 1. Kaplan-Meier plot of the probability of remaining free of tuberculosis according to body mass index (weight (kg)/height (m)²), First National Health and Nutrition Examination Survey (NHANES I) Epidemiologic Follow-up Study, 1971–1992.

8

Recent Studies 2021

THE LANCET
Infectious Diseases
Volume 21, Issue 10, October 2021, Pages e318–e325



Practical Topic

Food for thought: addressing undernutrition to end tuberculosis

Pranay Sinha MD^{1,2,3,4}, Kunt Linnroth PhD⁵, Anurag Bhargava MD^{6,7}, Scott K Heyesell MD⁸, Sonali Sarkar MD⁹, Padmini Salgame PhD¹⁰, William Rudgard PhD¹¹, Della Boccia PhD¹², Daniel Van Aartsen MD¹³, Natasha S Hochberg MD^{14,15}

Show more

Undernutrition is the leading population-level risk factor for tuberculosis.

Studies have consistently found that undernutrition is associated

- ✓ Increased tuberculosis incidence
- ✓ Increased severity
- ✓ Worse treatment outcomes
- ✓ Increased mortality

[Food for thought: addressing undernutrition to end tuberculosis - The Lancet Infectious Diseases](#)

9

RESEARCH ARTICLE

Effect of malnutrition on radiographic findings and mycobacterial burden in pulmonary tuberculosis

Kacie J. Hoyt^{1*}, Sonali Sarkar², Laura White³, Noyal Mariya Joseph⁴, Padmini Salgame⁵, Subitha Lakshminarayanan², Muthuraj Muthaiah⁶, Saka Vinod Kumar⁷, Jerrold J. Ellner⁸, Gautam Roy², C. Robert Horsburgh, Jr^{1,3,8}, Natasha S. Hochberg^{1,8*}


<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0214011>

Conclusion:

Malnutrition was associated with **increased extent of disease and cavitation on CXR**

10

**Nutritious
Food
Rest
Sunshine
Fresh Air**



THE CONSUMPTIVE
WORKING MAN.
What can Sanatoria do for Him?

BY
NORRIS DEAN HARKNELL
M.D., M.B., F.R.C. (Edin.)
PHYSICIAN IN CHARGE, WEST LONDON DISPENSARY

LONDON:
THE SCIENTIFIC PRESS LIMITED
41 & 43 DUCKWORTH STREET, DUCKWORTH, E.C.

1910.

CONCLUSIONS: VALUE OF SANATORIUM TREATMENT. 161

After Periods Varying from One to Five and a Half Years after Discharge.

Persons Treated.	In Normal Health and at Full Work.	Little or no Work, and in Poor or only Fair Health.	Death.
3	50 $\frac{0}{10}$	19-6 $\frac{0}{10}$	30-4 $\frac{0}{10}$
nd, 173	46-2 $\frac{0}{10}$	22-5 $\frac{0}{10}$	31-2 $\frac{0}{10}$
nd Sheffield,	41-7 $\frac{0}{10}$	16-6 $\frac{0}{10}$	41-7 $\frac{0}{10}$

BY
NORL GRAM HARDWELL
M.D., M.B., OF LONDON.
MEMBER OF THE ROYAL MEDICAL SOCIETY OF LONDON.


110109
HISTORICAL PAPER LAMING
HISTORICAL PAPER LAMING, N.C.

The actual Nutritive value of the daily diet was, protein, 196.5 grammes, fat 126.4 grammes and carbo-hydrate 522.6 grammes with a caloric value of 4.040"

6


Vitamin A May Help Boost Immune System to Fight Tuberculosis

Nutrient lowers intracellular cholesterol used by TB to sustain infection



UCLA Researchers

UCLA's Elliott Kim, Philip Liu and Avelino De Leon
February 25, 2014



13

MTB is Sensitive to Killing by a vitamin C-induced Fenton Reaction

VITAMIN C

Foods Sources

V	Valencia Orange	Cabbage, Red	Orange
I	Issai Kiwi Fruit	Cantaloupe	Papaya
T	Turnip Greens	Carambola	Pineapple
A	Apricots	Cauliflower	Potato
M	Mango	Cauliflower, Green	Prickly Pears
I	Ivy Gourd	Collard Greens	Pummelo
N	Nori	Chili Pepper, Hot	Radishes
C	Cantaloupe	Gooseberries	Raspberries
	Apricots	Grapefruit	Rutabagas
	Beans, Yellow Snap	Guavas	Spinach
	Bell Pepper	Kiwifruit	Squash, Summer
	Blackberries	Lemon	Strawberries
	Broccoli	Lime	Sweet Potato
	Brussels Sprouts	Nori	Tangerines
	Cabbage, Green	Mango	Tomato
	Cabbage, Pe-Tsai	Melon, Honeydew	Watermelon
		Okra	
		Onion	

On April 4, 1932 Vitamin C was first isolated by CC King at the University of Pittsburgh.



Department of Microbiology and Immunology, Howard Hughes Medical Institute NY, USA. 2013


14

Vitamin D Powerful Weapon Against TB

Researchers found that, in the presence of even minimally adequate levels of vitamin D, the body's own immune system will naturally trigger an immune response against the TB.

Journal Science Translational Medicine.

October 14, 2011



15

Most Recent Systematic Review

Effects of Vitamin D Supplementation on the Outcomes of Patients With Pulmonary Tuberculosis

A Systematic Review and Meta-Analysis


Hong-xia Wu; Xiao-feng Xiong; Min Zhu; Jia Wei; Kai-quan Zhuo; De-yun Cheng Disclosures BMC Pulm Med. 2018;18(108)

CONCLUSIONS:

Vitamin D supplementation can be considered as a combination therapy in patients with PTB.


16

Undernutrition and TB



17

Why is Nutrition Important in a Person with TB?



18

Importance of Nutrition in TB Treatment Response

Lack of Weight Gain & Relapse Risk in a Large Tuberculosis Treatment Trial

A. Khan, T. Sterling, R. Reeves, A. Vernon and the TB Trials consortium
American Journal of respiratory and Critical Care Medicine. Vol. 174

19

Importance of Nutrition in TB Treatment Response



HEARTland
NATIONAL TB CENTER
A PARTNERSHIP OF UT HEALTH SCIENCE CENTER AND TBCU

Impact of Poor Nutrition on TB Relapse

Reference: Lack of Weight Gain and Relapse Risk in a Large Tuberculosis Treatment Trial: Awa Khan, Timothy R. Sterling, Randall Reeves, Andrew Vernon, C. Robert Horsburgh and the Tuberculosis Trials Consortium; *American Journal of Respiratory and Critical Care Medicine* Vol 174, pp. 344-348, (2006)

Weight gain of 5% or less during the first 2 months of therapy is associated with an increased risk of relapse, even after controlling for other factors.

Consultation to healthcare providers at 1-800-TEX-LUNG
2303 SE Military Drive, San Antonio, TX 78223
www.HeartlandNTBC.org

This product produced with funds awarded by the Centers for Disease Control & Prevention (CDC)
Created 3-08

Khan Am J Resp Crit Care Med 2006

20

Importance of Nutrition in TB Treatment Response

- ✓ The relationship between nutritional status and poor outcomes for patients with TB.
- ✓ The association of weight gain between diagnosis and the end of 2-month Initial Phase therapy and risk of relapse

21

Definition of TB Relapse

Patients remain **culture negative** during treatment , **but after** completion of therapy, they become **culture positive** again or show clinical or radiographic deterioration consistent with active TB.

22

Lack of Weight Gain and Relapse Risk

- **857** subjects were enrolled.
- Monitored for two (2) years.
- **Body weight (kg) was measured at:**
 - ✓Diagnosis
 - ✓Enrollment in study
 - ✓Monthly during treatment
 - ✓And every 3-6 months during follow-up
- **Height**
- **BMI** (Body Mass Index)
- **IBW** (Ideal Body Weight)

Lack of Weight gain & Relapse Risk in a Large Tuberculosis Treatment Trial : A. Khan, T. Sterling, R. Reeves, A. Vernon and the TB Trials consortium . American Journal of respiratory and Critical Care Medicine. Vol. 174

23

WEIGHT AS A RISK FACTOR FOR TB RELAPSE

Underweight at Diagnosis ≥ 10% Below Ideal Body Weight			
	Weight gain after 2 months Rx	Relapse (%)	Cavitary AND Positive 2 months culture
Yes	≤ 5%	20.3%*	50.5%**
	> 5%	11.9%	18.5%
No		4.2%	18.3%

*p=0.06 **p=0.02

BMI	RELAPSE (5)
< 18.5	19.5%
18.51-19.0	10.7%
>19.0	6.1%

Body Mass Index (BMI) is optimal weight for health. Adults with a BMI between 19 and 24 have less risk for illnesses such as heart disease and diabetes than individuals with a BMI between 25 and 29. A BMI greater than 30 indicates greatest risk for obesity-related diseases. (See Chart 1.)

Adapted from The National Institute of Health, NHLBI Clinical Guidelines on Overweight and Obesity June 1998. www.nhlbi.nih.gov/guidelines.

24

Lack of Weight Gain and Relapse Risk

Results

61 patients relapsed (7.1%)

BMI	RELAPSE (5)
< 18.5	19.5%
18.51-19.0	10.7%
>19.0	6.1%

Khan. 2006 Am J Resp & Crit Care Med;174:34



25

Lack of Weight Gain and Relapse Risk

Underweight at Diagnosis ≥ 10% Below Ideal Body Weight			
	Weight gain after 2 months Rx	Relapse (%)	Cavitary AND Positive 2 months culture
Yes	≤ 5%	20.3%*	50.5%**
	> 5%	11.9%	18.5%
No		4.2%	18.3%

26

Remember...

Patients with **10% below ideal body weight**
at diagnosis **that don't regain at least 5%**
weight by end of two months of Rx



At 2 months
sputum
culture (+)

50% chance of relapse



27



28

Assessing Nutritional Status in a Person with TB



29

Laboratories (Normal Values)

Albumin: 3.8 – 5.2 g/dl
(Major protein. Low levels in poor diets, ↓ iron intake)

Total Protein: 6.0-8.5 g/dl (Low levels indicate poor nutrition)

Hemoglobin: 11.5 – 16 g/dl ♀ 13.2 – 17.1 g/dl ♂

Hematocrit: 36.0 – 45.0 % ♀ 38.5 – 50.5 % ♂


Glucose: 65 – 110 mg/dl

WBC: 3.8 – 10.8

Lymph: 18-48 % (decreases with progressive malnutrition)

30

Body Mass Index (BMI)



31

Ideal Body Weight Table

METROPOLITAN LIFE TABLE FOR MEDIUM FRAME ADULT

Ideal Weight For Women


Height in Shoes	Medium Frame
6'	148 to 162 lb
5'11"	145 to 159 lb
5'10"	142 to 156 lb
5'9"	139 to 153 lb
5'8"	136 to 150 lb
5'7"	133 to 147 lb
5'6"	130 to 144 lb
5'5"	127 to 141 lb
5'4"	124 to 138 lb
5'3"	121 to 135 lb
5'2"	118 to 132 lb
5'1"	115 to 129 lb
5'	113 to 126 lb
4'11"	111 to 123 lb
4'10"	109 to 121 lb

Ideal Weight For Men

Height in Shoes	Medium Frame
6'4"	171 to 187 lb
6'3"	167 to 182 lb
6'2"	164 to 178 lb
6'1"	160 to 174 lb
6'	157 to 170 lb
5'11"	154 to 166 lb
5'10"	151 to 163 lb
5'9"	148 to 160 lb
5'8"	145 to 157 lb
5'7"	142 to 154 lb
5'6"	139 to 151 lb
5'5"	137 to 148 lb
5'4"	135 to 145 lb
5'3"	133 to 143 lb
5'2"	131 to 141 lb


From height and weight tables of the Metropolitan Life Insurance Company, 1983. The ideal weights given in these tables are for ages 25 to 59. The weights assume you are wearing shoes with 1-inch heels and indoor clothing weighing 3 pounds.

32





Nutritional Teaching TIPS!

- ✓ Considerer Prolonging therapy for patients >10% underweight.
- ✓ Calculate BMI and IBW %
- ✓ Monitor weight weekly in underweight patients.
- ✓ Once stable, monitor monthly
- ✓ Ideally patients should gain 1lb/week
- ✓ Provide food resources
- ✓ Recommend iron-rich food intake if client is anemic
- ✓ Recommend intake of food sources of vit A, C, Vit D (fish, butter, milk etc)
- ✓ Encourage the patient to monitor his/her weight.



33

More Studies!


Int J Tuberc. Lung Dis. 2014 May;18(5):564-70. doi: 10.5588/ijtld.13.0602.

Body mass index predictive of sputum culture conversion among MDR-TB patients in Indonesia.

Compared to patients with **normal weight (BMI ≥18.5)**, **severely underweight patients (BMI <16)** had longer time to initial conversion and a lower probability of sputum culture conversion within 4 months.

Conclusion:
Severe underweight was associated with **longer time to initial sputum culture conversion** among MDR-TB patients.


34



35


A presentation slide with a white background and a blue decorative border on the right. On the left side, there is a circular logo featuring a stylized star or flower design. The title "Case Study" is centered at the top in a bold, dark blue font. Below the title is a photograph of a severely undernourished man standing in a bathroom. Below the photo is a list of symptoms in black text, with one item highlighted in red.

Case Study



- Chronic diarrhea, severe undernutrition, difficulty walking, generalized weakness
- **60Lb weight loss**
- Disseminated TB involving lungs and bowel

36



Nutritional Status:

Weight at admission: **77.8 Lb**

Height: **5'7"**

IBW (Ideal Body Weight): **142 Lb**

BMI : 12.2 Severely underweight


How to calculate the % IBW?

$$\% \text{ IBW} = \frac{\text{Current Body Weight}}{\text{Ideal Body Weight}} \times 100$$

$$\% \text{ IBW} = \frac{77.8}{142 \text{ lb.}} \times 100 = 54.7 \%$$

45.3 % ↓
Below of the IBW

37



Nutritional Update

Diet advance slowly
Patient refuses to eat meals on regular basis
After 1 year of treatment

Weight at d/c: **114 Lb**


Height: **5'7"**

IBW (Ideal Body Weight): **142 Lb**

BMI : 18 Underweight

$$\% \text{ IBW} = \frac{114 \text{ lb}}{142 \text{ lb.}} \times 100 = 80\%$$

38



39


Nutritional Status:

Weight at diagnosis: 91 Lb

Height: 5'2"

IBW (Ideal Body Weight): 123 Lb

BMI : 17 - Underweight



15-year-old Male

• < 3% on growth
• BMI 17

How to calculate the % IBW?


$$\% \text{ IBW} = \frac{\text{Current Body Weight}}{\text{Ideal Body Weight}} \times 100$$

$$\% \text{ IBW} = \frac{91 \text{ lb}}{123 \text{ lb}} \times 100 = 73.9\%$$

26.2% ↓


Below of the IBW

40



Nutritional Update

- Patient was treated for PTB for 6 months (non-cavitary)
- Episode of neutropenia
- Clinical improvement Increase energy, appetite and
- **Gained 15 lb.**



Weight at en of Rx: 106 Lb


Height: 5'2'

IBW (Ideal Body Weight): 123 Lb

BMI : 19.2 Normal weight

$$\% \text{ IBW} = \frac{105}{123 \text{ lb.}} \times 100 = 85\%$$

41



**"Giving people medicine for TB
and not giving them food is like
washing your hands and drying
them in the dirt"**

Quote by a Haitian public health worker
Book: Mountains Beyond Mountains

42

42

