



# **Overview of Contact Identification Guidelines**

Matthew D. Whitson, RN  
September 18, 2024

Introduction to TB Nurse Case Management Online  
September 4<sup>th</sup> – September 25<sup>th</sup>, 2024  
Online Course

**Matthew D. Whitson, RN** has the following disclosures to make:

---

- No conflict of interests
- No relevant financial relationships with any commercial companies pertaining to this educational activity





TEXAS  
Health and Human  
Services

Texas Department of State  
Health Services

# Overview of Contact Identification Guidelines

---

**Matthew D. Whitson, RN**

**TB Nurse Consultant**

**Department of State Health Services, Region 8**

# Objective



TEXAS  
Health and Human  
Services

Texas Department of State  
Health Services

- Discuss the impact of contact investigation guidelines on existing contact investigation procedure.
- Links to document:

<https://www.cdc.gov/mmwr/pdf/rr/rr5415a1.pdf> (.pdf)

<https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5415a1.htm> (.htm)





**TEXAS**  
Health and Human  
Services

---

**Texas Department of State  
Health Services**

# **Why do we perform contact investigations?**

---

# Example #1

---

- 21 year old contact to a patient with TB disease. Tested QFT positive when initially evaluated and was non-adherent with all follow-up requests from the health department.
- One year later (22 years old), patient is diagnosed with active TB disease.
  - Patient has 1 y.o. child at home requiring window treatment.
  - Patient babysits a 3 y.o. child, also requiring window treatment.
  - Patient was enrolled in school at the time, requiring testing of school contacts.
  - Patient continues to have ongoing side effects to her medication regimen.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# A hopeful start

---

“Contact investigations are complicated undertakings that typically require hundreds of interdependent decisions, the majority of which are made on the basis of incomplete data, and dozens of time consuming interventions.” p. 1

Excerpt from Guidelines for the Investigation of Contacts of Persons with Infectious Tuberculosis, 2005



TEXAS  
Health and Human Services

Texas Department of State  
Health Services



# Methodology



TEXAS  
Health and Human  
Services

Texas Department of State  
Health Services

- Based upon on epidemiologic and other relevant scientific studies and established practices in conducting CIs.
- CI has not been researched by a controlled trial or study, but through expert opinion from years of common practices.
- These guidelines are not “one size fits all”.





# Decision to Initiate Contact Identification

---

## General issues

- Competing Demands
  - Limited resources that may be allocated
  - Limited staff
- Prioritizing Cases
  - Which cases are the most and least likely to involve transmission of disease?
  - Which cases have the highest priority contacts?
  - Which cases are the highest profile?



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Decision to Initiate Contact Identification

---

## TB transmission factors

- Only patients with **pulmonary, laryngeal** or **pleural TB** can transmit their infection (*with few exceptions*).
  - In some rare cases, extrapulmonary disease can cause transmission during medical procedures that release aerosols (autopsy, embalming, and irrigation of a draining abscess).
- Extrapulmonary disease cases should not be given high priority unless accompanied by pulmonary, laryngeal or pleural disease.



TEXAS

Health and Human Services

Texas Department of State  
Health Services

# Decision to Initiate Contact Identification

## TB transmission factors

- Anatomical site of disease
- Sputum bacteriology
- Radiographic findings
- Behaviors that increase aerosolization of respiratory secretions
- Age
- HIV status
- Administration of effective treatment





FIGURE 1. Decision to initiate a tuberculosis (TB) contact investigation



\* Acid-fast bacilli.

† Nucleic acid assay.

§ According to CDC guidelines.

‡ Chest radiograph.

# Interviewing patients

---

## Patient interview

- Health department's responsibility to conduct CI.
  - Clearly state written policies and procedures.
  - Improves efficiency and uniformity.
- Establishing trust and consistent rapport with patients is critical.
  - Gain full information and cooperation during treatment.
- Interviews should be in primary language of the interviewee.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Interviewing patients

## Determining Infectious Period

- Infectious period
  - **3 months** before a TB diagnosis
  - 4 weeks before possible diagnosis if no symptoms, no positive sputum, no cavities
- In certain circumstances, an even earlier start date should be used.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services





Texas Department of State  
Health Services

TABLE 2. Guidelines for estimating the beginning of the period of infectiousness of persons with tuberculosis (TB), by Index case characteristic

TB symptoms	Characteristic		Recommended minimum beginning of likely period of infectiousness
	AFB* sputum smear positive	Cavitary chest radiograph	
Yes	No	No	3 months before symptom onset or first positive finding (e.g., abnormal chest radiograph) consistent with TB disease, whichever is longer
Yes	Yes	Yes	3 months before symptom onset or first positive finding consistent with TB disease, whichever is longer
No	No	No	4 weeks before date of suspected diagnosis
No	Yes	Yes	3 months before first positive finding consistent with TB

**SOURCE:** California Department of Health Services Tuberculosis Control Branch; California Tuberculosis Controllers Association. Contact investigation guidelines. Berkeley, CA: California Department of Health Services; 1998.

\* Acid-fast bacilli.

# Interviewing patients

---

## Determining Infectious Period

- Stringent criteria should be applied when susceptible contacts are involved:
  - At least 3 consecutive negative sputum AFB smear results from sputum collected  $\geq 8$  hours apart (with one specimen collected during the early morning)



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Interviewing patients

## Closing Infectious Period

- Infectious period is closed when:
  - Effective treatment for  **$\geq 2$  weeks**
    - Sometimes after 5 days of treatment if AFB smears are all negative
- Diminished symptoms
- Mycobacteriologic response

AFB smears are trending downward or are negative



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Interviewing patients

---

## Things to remember

- Interview multiple times.
- Interview at patient's residence at least once. (Look for clues of other people.)
- Obtain medical records and information regarding the patient's illness.
- Conduct at least one interview (preferably the first) in person or with a proxy.
- Take the environment into consideration.
- Finding those exposed to TB is an ongoing process.



TEXAS

Health and Human Services

Texas Department of State  
Health Services

# Prioritization and Evaluation

---

## Assigning priorities

- Most recently infected contacts cannot be distinguished right away.
- Create a line list and ensure that every person has a break-in-contact (BIC) date.
- Repeat testing done 8-10 weeks after the BIC will help to identify recent conversions that are most likely indicative of recent infection.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Prioritization and Evaluation



## How to categorize

- Likelihood of infection depends on:
  - Intensity
    - How many germs was the person exposed to?
  - Frequency
    - How often was the person around the patient?
  - Duration of exposure
    - How long was the person around the patient?
  - Risk factors
    - Does the person have any age or medical risk factors that would place him or her at a greater risk of infection or disease?




TEXAS  
Health and Human Services

Texas Department of State  
Health Services



FIGURE 2. Prioritization of contacts exposed to persons with acid-fast bacilli (AFB) sputum smear-positive or cavitary tuberculosis (TB) cases



\* Human immunodeficiency virus or other medical risk factor.

† Bronchoscopy, sputum induction, or autopsy.

§ Exposure exceeds duration/environment limits per unit time established by the health department for high-priority contacts.

¶ Exposure exceeds duration/environment limits per unit time established by the health department for medium-priority contacts.

Prior

Sub

- 1 c



TEXAS  
Health and Human Services

Texas Department of State  
Health Services


10/3/2022

# Prioritization and Evaluation

FIGURE 3. Priority assignments for contacts exposed to persons with acid-fast bacilli (AFB) sputum smear-negative tuberculosis (TB) cases

Sub

- 1 c



\* Nucleic acid assay.

† Human immunodeficiency virus or other medical risk factor.

§ Bronchoscopy, sputum induction, or autopsy.

¶ Exposure exceeds duration/environment limits per unit time established by local TB control program for medium-priority contacts.




TEXAS  
Health and Human Services

Texas Department of State  
Health Services



TEXAS  
Health and Human Services  
Texas Department of State  
Health Services

**FIGURE 4. Prioritization of contacts exposed to persons with suspected tuberculosis (TB) cases with abnormal chest radiographs not consistent with TB disease**



\* Acid-fast bacilli.

† Nucleic acid assay.

§ Human immunodeficiency virus infection or other medical risk factor.

¶ Bronchoscopy, sputum induction, or autopsy.




TEXAS

Health and Human Services

Texas Department of State  
Health Services

Low -  
medium  
risk



Work

Family

Family

Work

Social

Social

Calculate positivity  
rates and move  
outward accordingly

High risk



# Prioritization and Evaluation

---

## How to evaluate contacts

- Initial encounter with high- and medium- priority contacts who are most at risk should be made **within 3 working days.**
  - In Texas, the form is known as TB-340.
  - Gather contact's background health information.
  - Face-to-face assessment of the person's health.
  - Draw IGRA or administer TST at that time.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Prioritization and Evaluation


## How to evaluate contacts

- The health department's evaluation for TB or LTBI should include:
  - Previous M.TB infection or disease and related treatment.
  - Contact's verbal report and documentation of previous TST results.
  - Current symptoms of TB illness
  - Risk factors or medical conditions making TB disease more likely
  - Mental health disorders
  - Type, duration, and intensity of TB exposure
  - Sociodemographic factors



TEXAS  
Health and Human Services

Texas Department of State  
Health Services





# Prioritization and Evaluation

---

## How to evaluate contacts

- Test for infection using IGRA or TST.
  - Stick with one test if it is valid; don't do best 2 out of 3 results.
- For children less than 5 years of age, have HIV, or patients who are immunocompromised, perform a PA and lateral CXR.
- Evaluate for signs and symptoms of TB disease.
- Patient must be evaluated again 8-10 weeks after the BIC.




TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Prioritization and Evaluation

## IGRA<sub>s</sub>

- IGRA
  - QuantiFERON
  - T-spot
- Pros
  - One visit
  - Objective
  - Rules out false positives
- Cons
  - Expensive
  - Indeterminates, indicator of disease vs. infection
  - Immunosuppressed patients




TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Prioritization and Evaluation

## Further medical evaluation

- Recent Contacts whose skin test induration diameter is  $\geq 5$  mm or who report any symptoms consistent with TB disease:
  - Should undergo further examination and diagnostic testing for TB, starting typically with a chest radiograph.
  - Collection of specimens for mycobacteriologic testing is decided on a case-by-case basis and is not recommended for healthy contacts with normal chest radiographs.
  - All contacts who are classified as high priority due to risk factors or vulnerability should undergo further examination and diagnostic testing regardless of TST result or symptoms.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Prioritization and Evaluation

---

## Follow-up

- Documentation of a previous positive result should be obtained before omitting the skin test from the diagnostic evaluation.
- Do not make a decision whether or not to test based on a contact's verbal report.
- Provide window prophylaxis for high risk groups (< 5 y.o., HIV+, immunosuppressed)
- Determine if legal action is required for those who refuse testing for their children.




**TEXAS**  
Health and Human Services

Texas Department of State  
Health Services

# Data Management and Communications

## Collecting Data

- Collect specific data needed for evaluation.
- Collect on standardized forms
- Use specified standardized data definitions and formats when possible.
- Have a plan for how you are going to collect, process, and analyze data from the beginning.




TEXAS  
Health and Human Services

Texas Department of State  
Health Services

# Special settings and source case investigations

## Special settings

- Congregate or unusual settings for TB exposures
  - Correctional facilities
  - Workplaces
  - Hospitals and other healthcare settings
  - Schools
  - Shelters
  - Modes of Transportation
  - Drug and alcohol usage sites
  - Interjurisdictional investigations





# Special settings and source case investigations

---

## Outbreaks

- Definition with regard to TB:
  - During a CI, 2 or more contacts having TB
  - Any 2 or more cases occurring less than 1 year of each other are linked and not caught in a CI.
- May indicate lapse in regular TB control.
- Call for assistance when needed.
- The public may use the term “outbreak”, but a TB outbreak rarely occurs.



TEXAS  
Health and Human Services  
Texas Department of State  
Health Services

# Special settings and source case investigations

## Source case investigations

- Reverse contact identification
  - Goal is to actively search for someone who has TB disease.
  - Recommended for children under the age of 5 years old who develop TB disease.
  - Start close and work your way out.
  - Searching for unexplained TBI is generally not recommended and if done should be limited to LTBI in children younger than 2 years of age.



TEXAS  
Health and Human Services

Texas Department of State  
Health Services





# Take away points

- Maintain control
- Communicate effectively
- Stay organized
- Be flexible
- Always improve



TEXAS  
Health and Human  
Services

Texas Department of State  
Health Services





TEXAS  
Health and Human  
Services

Texas Department of State  
Health Services

# Thank you

Matthew D. Whitson, RN

[matthew.whitson@dshs.texas.gov](mailto:matthew.whitson@dshs.texas.gov)

210-949-2000